

Corporate Presentation

November 2022

Disclaimers

Important information

The Company is not making any offers of securities at this time and cannot accept any orders for securities at this time. This presentation shall not constitute an offer to sell or the solicitation of an offer to buy any securities of the Company.

The presentation does not purport to give legal, tax or financial advice, and must not be relied upon for any such purpose. Neither the information contained in this presentation, nor any further information made available by the Company or any of its affiliates or employees, directors, representatives, officers, agents, or advisors, in connection with this presentation will form the basis of or be construed as a contract or any other legal obligation.

This presentation may contain certain forward-looking statements and opinions. Forward-looking statements are statements that do not relate to historical facts and events and such statements and opinions. pertaining to the future that, by example, contain wording such as "believes", "estimates", "anticipates", "expects", "forecasts", "intends", "could", "will", "should", "would", "according to estimates", "is of the opinion", "may", "plans", "potential", "predicts", "to the knowledge of" or similar expressions, which are intended to identify a statement as forward-looking. This applies, in particular, to statements and opinions in this presentation concerning our future business expectations, continued FDA approval for TARPEYO, market acceptance of TARPEYO, clinical trials, supply chain, strategy, goals and anticipated timelines, competition from other biopharmaceutical companies, our future financial returns, plans and expectations with respect to our business and management, future growth and profitability and general economic and regulatory environment and other matters affecting our company. Forward-looking statements are based on current estimates and assumptions made according to the best of our knowledge. Such forward-looking statements are subject to risks, uncertainties, and other factors that could cause the actual results, including with respect to our business, cash flow, financial condition and results of operations, to differ materially from the results, or fail to meet expectations expressly or implicitly assumed or described in those statements or to turn out to be less favorable than the results expressly or implicitly assumed or described in those statements. Accordingly, you should not place undue reliance on the forward-looking statements herein. We can give no assurance regarding the future accuracy of the opinions set forth herein or as to the actual occurrence of any predicted developments. In light of the risks, uncertainties and assumptions associated with forward-looking statements, it is possible that the future events mentioned in this presentation may not occur. Actual results, performance or events may differ materially from those in such statements due to, without limitation, risks generally associated with pharmaceutical development, including delays or challenges that may arise in the clinical development of our product candidates, interactions with regulatory authorities, challenges in the commercialization of our product candidates, if approved, the risk that we may not maintain our existing collaboration or enter into new ones, or that we will not realize the intended benefits from such collaborations, any inability to expand our pipeline of product candidates, any inability to maintain barriers to entry such as intellectual property protection or regulatory exclusivity, changes in general economic conditions, in particular economic conditions in the markets on which we operate, changes affecting interest rate levels, changes affecting currency exchange rates, changes in competition levels and changes in laws and regulations. The information, opinions and forward-looking statements contained in this announcement speak only as at its date, and are subject to change without notice.

Certain information, including certain forward-looking estimates and forecasts, contained in this presentation relates to or is based on studies, publications, surveys and other data obtained from third-party sources and the Company's own internal estimates and research. While the Company believes these third-party sources to be reliable as of the date of this presentation, it has not independently verified, and makes no representation as to the adequacy, fairness, accuracy or completeness of any information obtained from third-party sources. In addition, all of the market data included in this presentation involves a number of assumptions and limitations, and there can be no guarantee as to the accuracy or reliability of such assumptions. Finally, while we believe our own internal research is reliable, such research has not been verified by any independent source.






A commercial stage biopharma company focused on novel treatments in orphan indications, with an initial focus on renal and hepatic diseases with significant unmet needs.

- Pioneers in immunoglobulin A nephropathy (IgAN);
 - Positive Phase 2b results published in The Lancet
 - Successful top-line data read out from the ongoing Phase 3
 NeflgArd trial in November 2020.
 - Both clinical trials met primary and key secondary endpoints.
 - December, 2021; the first and only FDA approved medication for this rare disease
 - July 2022, EC approval in Europe; Sept 2022, transfer of MA to partner STADA, who will launch in Germany in Q4 2022
- Strong pipeline in orphan liver and kidney indications; platform of first-in-class NOX Inhibitors.
 - Actively expanding through in-licensing and/or acquisition of product candidates.





Pipeline





First and Only Approved Medication for IgAN in Europe & USA





The first and only FDA approved drug specifically designed for Immunoglobulin A Nephropathy (IgAN)
Approved by FDA in December 2021; Launched in Jan 2022

The first and only EMA approved drug specifically designed for IgAN Positive CHMP Opinion in May 2022, European Commission approval in July 2022 Commercial Partner, STADA, received MA in September 2022 and will launch in Germany in Q4 2022

Launched under an accelerated approval to reduce proteinuria in adults with IgAN at risk of rapid disease progression, generally a urine protein-to-creatinine ratio (UPCR) ≥ 1.5 g/g



IgA Nephropathy – a significant unmet medical need


Profile

 Genetic predisposition is required but not sufficient; environmental, bacterial, dietary factors

may play a role

More than 50% are at risk of developing ESRD within 10-20 years, which can only be treated via regular haemodialysis or kidney transplant

 High levels of proteinuria is connected to disease progression and worse outcomes for patients



Estimated Prevalence

Market Opportunity



130,000 - 150,000



200,000




~5,000,000



Pathophysiology of IgAN – predominant theory

Patients with IgAN have an increased appearance in the blood of secretory IgA antibodies, which are produced in the Peyer's patches, that lack galactose units in the hinge region

These galactose-deficient IgA antibodies are immunogenic, triggering IgA and IgG autoantibody production directed against the hinge region



The galactosedeficient secretory IgA antibodies form immune complexes with the IgA and IgG autoantibodies


These immune complexes are deposited in the glomeruli of the kidney, causing an inflammatory cascade and destroying the glomeruli. This reduces the kidney's ability to remove waste products from the blood and eventually may result in ESRD



Designed to target the presumed origin of the disease


Drug product based on known active ingredient

- Active ingredient is budesonide a highly potent, locally acting corticosteroid
- 90% cleared in first pass metabolism by liver, with the view of minimizing systemic side effects
- Safety profile as expected for oral administration; predominantly mild to moderate AEs, reversible upon discontinuation



Novel targeted release profile

- Designed to deliver a targeted and highly potent dose directly to Peyer's patches in the ileum
- Differentiated release profile
 - pH-governed delayed disintegration of the capsule until it reaches the ileum
 - Potent, sustained exposure throughout the ileum






Positive Phase 3 Data – Part A Efficacy Results

First successful readout of a Phase 3 trial in IgA nephropathy

9 months of dosing with 16mg TARPEYO in 199 patients demonstrated a statistically significant and clinically meaningful reduction in proteinuria and in eGFR stabilization.



Efficacy Findings

- Statistically significant UPCR reduction with TARPEYO (16 mg) compared to placebo following 9 months treatment (p=0.0001)
- After 9 months
 - Reduction in UPCR for TARPEYO treated patients = 31%
 In ITT population, this reduction was 34%
 - Reduction in UPCR for placebo = 5%
- After 12 months
 - Reduction in UPCR for TARPEYO treated patients = 52%
 - Reduction in UPCR for placebo = 7%
- Statistically significant eGFR stabilisation After 9 months:
 - eGFR decline for TARPEYO treated patients = 0.17ml/min/1.73m²
 - eGFR decline for placebo = 4.04ml/min/1.73m²

Safety Findings

- Generally well-tolerated; majority of AR mild/moderate in severity
- No adverse clinical effects on the cardiovascular or metabolic system
 No severe infections



TARPEYO: A differentiated and targeted approach for IgAN

Targeted immunomodulator down regulating IgA1 at the source

Proteinuria reduction vs SOC of physicians choice significant continued reduction off drug

Designed to locally target origin of disease with the potential of being disease modifying

Local action complemented by limited systemic exposure due to first pass metabolism of 90%

Patient centric

Well characterized active ingredient and safety profile

Option of intermittent, rather than chronic treatment

Demonstrated impact not just on proteinuria but on eGFR in patients at risk of rapid progression





TARPEYO: Defining the market with the first and only FDA approved drug in IgAN

TARPEYO[™] (budesonide) delayed release capsules is a corticosteroid indicated to reduce proteinuria in adults with primary immunoglobulin A nephropathy (IgAN) at risk of rapid disease progression, generally a urine protein-to-creatinine ratio (UPCR) \geq 1.5 g/g. This indication is approved under accelerated approval based on a reduction in proteinuria. It has not been established whether TARPEYO slows kidney function decline in patients with IgAN. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory clinical trial.





Established highly successful support service for frictionless access



- TARPEYO TouchpointsTM: full-service patient and provider support program. Fully operational on day 1 of TARPEYO approval
- Utilizes Biologics by McKesson's PharmacyEliteTM model; integrated HUB* and exclusive Specialty Pharmacy
- Staffed by Care Navigators: dedicated case managers + designated Rare Pod Team (nurses, pharmacists, fulfillment and distribution team)
- Integrated with a financial assistance (commercial co-pay) program provided by CoverMyMeds® from McKesson



Our US Commercial Launch leadership team of industry experts



Extensive launch expertise: commercial experience at top-tier pharma (eg, Pfizer, Bayer, BMS, Regeneron)




Experienced sales force: 60 sales reps with core background in rare disease, specialty products and nephrology market



Hands-on managers: 6 national account managers in the field and engaging targeted payers




Expert partners: AmerisourceBergen (ICS), McKesson (Biologics), CDM and LifeSci Communications





TARPEYO Launch Metrics





Total Unique Prescribers: 480

Total Enrolments: 730



Continued growth through Q3 2022

- Strong net sales
- Continued penetration of prescriber base
- Encouraging and broad interest from nephrology community
- Market Access reached goal of more than 90% of US lives covered
- Payer mix less than 25% government insured with **remainder either private insurance or cash**
- Awareness continues to grow now over 80%
- Advocacy and patient meetings and activities ongoing
- Clinical data recently published in peer reviewed journal





Ex-US Commercial Partnerships



2019: Partnered with Everest Medicines in Greater China and Singapore

Initial upfront payment of 15M USD Up to an additional 106M USD in future milestone payments, as well as royalties on annual net sales at a low- to midteens percentage

Strong expertise in clinical development and regulatory arena with an innovative biopharm approach. Nine other products in the pipeline, 3 others FDA-approved

Public company listed on the Hong Kong exchange (1952.HK)



2021: Partnered with STADA in the European Economic Area (EEA) member states, Switzerland, and the UK

Initial upfront payment of 20M EUR (\$20M) and up to an additional 77.5M EUR (\$77.6M) in future milestone payments, as well as tiered royalties on net sales at a low 20s to low 30s percentage. \$13M EUR (\$13M) milestone payment following European Commission approval in July 2022

Extensive marketing and sales platform across Europe; large team with over 12,400 FTE

Private company with reported revenue in 2021 of 3,249.5M EUR (\$3,256M), EBITDA of EUR 776.5M EUR (\$778.1M)








NOX Inhibitor Development Program

A first-in-class platform

NOX Inhibitors

- Calliditas' development programs are based on a first in class, novel NOX inhibitor platform that includes lead compound setanaxib, the first NOX inhibitor to reach the clinical trial stage.
- NOX enzymes are solely dedicated to producing reactive oxygen species (ROS) as their primary and sole function. There are seven NOX members, each differing in composition, modes of activation and the ROS type they produce.
 - NOX1, NOX2, NOX3, and NOX5 transfer electrons from NADPH to molecular oxygen, producing superoxide anion (O_2^{-}) . NOX4, DUOX1 and DUOX2, meanwhile, mainly produce hydrogen peroxide (H_2O_2) .




At appropriate concentrations, ROS have essential functions in cellular signalling processes, helping to regulate cell proliferation, differentiation and migration, as well as modulating the innate immune response, inflammation and fibrosis. However, disruption of redox homeostasis has been implicated in multiple disease pathways.




Setanaxib in Squamous Cell Carcinoma of the Head & Neck

Scientific rationale

- The response to immuno-oncology therapies can be affected by the tumour microenvironment, in particular by the numbers of tumour-infiltrating lymphocytes (TILs) and cancer-associated fibroblasts (CAFs) in the tumour.
- A relationship between cancer associated fibroblasts (CAFs) and prognosis in squamous cell carcinoma of the head & neck (SCCHN) has been established.
- NOX4 is highly over-expressed in CAFs and drives myofibroblastic activation within tumours, shielding them from CD8+ TILs. Targeting CAFs with setanaxib could improve patients' responses to immunotherapies, and function as an adjunctive.
- There is increasing use of pembrolizumab as 1st line monotherapy in patients with relapsed or metastatic SCCHN, although response rates are low (ORR approx. 20%).









Preclinical work: Combining GKT137831 with aPD-1 in CAF-rich tumours (TC1)

- Using a CAF-rich tumour model in mice, administration of setanaxib + pembrolizumab (versus either treatment alone) resulted in:
 - -Improved penetration of TILs into the centre of the tumour
 - -Slowing of tumour growth
 - Improved survival





Setanaxib Phase 2b/3 (TRANSFORM) Trial in Primary Biliary Cholangitis (PBC)

- Double-blind, randomized, placebo-controlled, adaptive study design
 - Primary endpoint: ALP <1.67x ULN, and ALP reduction >15%, and total bilirubin <ULN
- ~318 Patients with PBC, elevated liver stiffness, and inadequate biochemical response/ intolerance to UDCA
 - 52 weeks of treatment with setanaxib 1200mg daily, setanaxib 1600mg daily, or placebo
 - Interim/ futility analysis after 99th patient has completed 24 weeks of treatment; will determine what dose of setanaxib to select for Phase 3
 - A 52 week extension phase, where all patients will receive blinded treatment with setanaxib
- Interim analysis targeted for 1H 2024, subject to recruitment
- Calliditas received FDA fast track designation for setanaxib in PBC in August 2021



Anticipated Milestones

Anticipated milestones regarding Calliditas' clinical, regulatory and commercial plans

2H 2021

- ➤ European commercial partnership ✓
- ➤ FDA target PDUFA date accelerated approval for TARPEYO in IgAN
- ➤ Initiate pivotal
 Phase 2/3 TRANSFORM
 trial in PBC ✓

2022

- ➤ Commercial launch of TARPEYO in the US <
- ➤ Initiate proof-of-concept Phase 2 trial in head and neck cancer ✓
- ➤ Positive EMA opinion for conditional approval for Kinpeygo (Nefecon) and European Commission Approval ✓
- ➤ Commercial launch in Europe (Q4)
- Regulatory filing in China (Q4)
- Commercial ramp in US

2023

- Readout of NeflgArd study Part B; completion of Phase 3 program
- > Filing for full approval in primary IgAN
- ➤ Data from head and neck cancer trial, subject to recruitment
- ➤ Potential approval in China
- ➤ Commercial ramp in EU

